

This report card summarises the
2022 Native Fish Report Card
(NFRC) survey in the Glenelg River.

SITES: 10


ELECTROFISHING

Fish found in the Glenelg River for NFRC



Target Species

✓ recorded in 2022



✓ **Estuary Perch**

Percalates colonorum



Non-target species

✓ recorded since 2017*

Large-bodied native species

- ✓ Australian Bass
- ✓ Australian Grayling
- ✓ Black Bream
- ✓ Freshwater Catfish
- ✓ Golden Perch
- ✓ River Blackfish
- ✓ Short-finned Eel
- ✓ Tupong
- ✓ Yellow-eye Mullet

Small-bodied native species

- ✓ Australian Smelt
- ✓ Carp Gudgeon
- ✓ Common Galaxias
- ✓ Flatheaded Gudgeon
- ✓ Southern Pygmy Perch
- ✓ Variegated Pygmy Perch

Exotic species

- ✓ Common Carp, Eastern Gambusia, Goldfish, Rainbow Trout, Redfin, Tench

* Incidentally captured but not measured as for target species.

Fish community

The NFRC Program began in 2017, with a focus on targeting the monitoring of population dynamics of key iconic fish species that have high recreational and/or conservation values, in large rivers across Victoria. In the Glenelg River, the target species is Estuary Perch. Surveys occur in February/March each year, at 10 sites from Dartmoor to Yat Nat (between Balmoral and Rocklands Reservoir). The equipment and habitats surveyed are focused on this species, which are measured to determine population structures. Other fish species that are incidentally captured are recorded, but not measured to determine their population structures.

Summary of key health indicators for target species in 2022

Species	Key Health Indicators		
	Recent recruitment	Multiple size classes	Mature fish present
Estuary Perch	No	Yes	Yes

Recent recruitment means young-of-year fish

Estuary Perch are considered an estuarine species, which often moves into lower freshwater reaches of rivers, particularly rivers with elevated baseline salinity levels.

Non-target species

The non-target fish species that have been incidentally recorded in the Glenelg River during NFRC surveys since 2017 are:

Large-bodied native species

Other large-bodied species recorded in surveys are Australian Bass, Australian Grayling, Black Bream, Freshwater Catfish, Golden Perch, River Blackfish, Short-finned Eel, Tupong and Yellow-eye Mullet. A single Australian Grayling was collected in 2019 with another single individual recorded in 2021. These are only the second and third records of this species in the Glenelg River system; the other confirmed record was 124 years ago. Black Bream and Yellow-eye Mullet are estuarine species that can move upstream into lower freshwater reaches of streams. River Blackfish are a lowland species, generally found at altitudes below 200 metres. This species has suffered a decline in distribution and abundance across the State¹, however, numerous individuals have been detected in the Glenelg River (ARI unpublished data). Short-finned Eel are a diadromous (migratory between salt water and fresh water) species found throughout coastal Victoria.

Large-bodied native species which have been introduced to the river

Australian Bass, Freshwater Catfish and Golden Perch are considered translocated species in the Glenelg River.

Small-bodied native species

Some of the small-bodied species recorded within the Glenelg River, including Australian Smelt and Flatheaded Gudgeon are common across the state. Carp Gudgeon are a lowland species and considered often hard to detect via boat electrofishing. The Common Galaxias and Tupong are diadromous species common across coastal Victoria. Pygmy Perch species are more common in offstream habitats such as billabongs, wetlands and lagoons. Variegated Pygmy Perch have been detected in all five years of NFRC sampling.

Exotic fish species

Common Carp, Eastern Gambusia, Goldfish and Redfin are widely distributed across sampling sites, but the likelihood of their detection and abundances increases as you move upstream. Small (young-of-year) Carp were also detected at most sites in 2019. In 2017-18 and 2020-22, small Carp have been restricted to the upper sites, indicating the widespread dispersal and spawning of this species during the 2018 spawning season. The presence of small Carp every year at the upper sites also indicates they are spawning in these areas in most, if not all years. Rainbow Trout are restricted in distribution to the Warrock area. Tench have been detected in all six sampling years, albeit in low abundances.

Other native fish species known from the Glenelg River

Some fish species known to occur in the Glenelg River have never been recorded during NFRC surveys. For example, no Climbing Galaxias, Obscure Galaxias, Spotted Galaxias, Little Galaxias, Pouched Lamprey or Short-headed Lamprey have been detected in the surveys. Both Climbing Galaxias and Spotted Galaxias historically had patchy distributions within the Glenelg River system and are hard to detect using the NFRC sampling methodology. Similarly, both lamprey species had patchy distributions historically, while Obscure Galaxias are hard to detect using the NFRC sampling methodology. Little Galaxias are normally found in lower altitude areas but are found in the Glenelg River upstream Rocklands Reservoir (outside of the NFRC sampling area).

Other notable species

Surveys have also recorded Eastern Long-necked Turtles, Platypus and Yabbies.

Environmental and Management Context

Environment

Low flow conditions were present in all six sampling seasons. Two sites were not fished in 2022 due to limitations in access.

River rehabilitation efforts in the Glenelg River

Many rehabilitation actions have occurred, and are underway, to improve the health of the Glenelg River. These are informed by the Glenelg Hopkins Waterway Strategy 2014-2022 and the Glenelg River Restoration Program. Actions include revegetation, weed control and fencing of riparian areas, reintroduction of instream woody habitat, allocations of water for the environment and removal of migration barriers and pest control. There are a range of fish monitoring efforts related to the rehabilitation efforts. These include the Victorian Environmental Flow Monitoring and Assessment Program ([VEFMAP](#)). The [Glenelg Hopkins Catchment Management Authority](#), DELWP and the [Victorian Fisheries Authority](#) support rehabilitation and management of the Glenelg River and its fish community.

See the ARI website for more information on the [Native Fish Report Card program](#).

¹ Khan et al. (2004). Habitat use and movement of river blackfish (*Gadopsis marmoratus* R.) in a highly modified Victorian stream, Australia. *Ecology of Freshwater Fish*, 13: 285–293.

The NFRC program, and related monitoring initiatives, provide improved understanding of the structure of fish communities and how rivers can be best managed.




Figure 1. Map showing the section of Glenelg River where NFRC sampling occurs




Figure 2. A range of size classes of Estuary Perch have been recorded during NFRC surveys



Figure 3. An adult Estuary Perch

Estuary Perch

Percalates colonorum



Key Health Indicators

- ✗ Recent recruitment
- ✓ Multiple size classes
- ✓ Mature fish present

Monitoring Results

Total number of fish caught	18
Fish per 1km of waterway	2.79
Largest fish by length (cm)	45.1
Largest fish by weight (kg)	1.26
% of the catch that is legal size	33.3

GLENELG RIVER

RECREATIONAL SPECIES

Estuary Perch (*Percalates colonorum*) - formerly *Macquaria colonorum* - are an estuarine species that can push into lower freshwater reaches of streams. Within the Glenelg River, the species pushes further inland than other rivers in Victoria. No Estuary Perch have been detected as far upstream as Balmoral, but sites have been included up this far, as the species seems to be slowly moving further upstream (based on previous surveys). Abundance of Estuary Perch was relatively consistent between all six years, though they appear to be declining, with the lowest number of fish per kilometre recorded in 2022 (Figure 4). Any interpretation of this result should however note that two sites where fish would normally be caught could not be surveyed in 2022 since they were inaccessible. Multiple size classes including mature and juvenile fish were captured in all years, with young-of-year detected in 2018. This is an indication that conditions in the Glenelg River are supporting spawning and survival of this species throughout its lifecycle and providing good, ongoing angling opportunities. A wide range of sizes was collected in 2022, but the 20-25 cm size range dominated the catch (Figure 5).

Stocking

No stocking has occurred.




Figure 4. The densities of recruits, juveniles and adult Estuary Perch for NFRC surveys in the Glenelg River from 2017 to 2022




Figure 5. The size range percentage of Estuary Perch measured from the Glenelg River during NFRC surveys in 2022

We acknowledge and respect Victorian Traditional Owners as the original custodians of Victoria's land and waters, their unique ability to care for Country and deep spiritual connection to it. We honour Elders past and present whose knowledge and wisdom has ensured the continuation of culture and traditional practices.